

IOWA STATE UNIVERSITY

Digital Repository

Swine Research Report, 2001

Animal Science Research Reports

2002

Analysis of National Animal Health Monitoring System Tonsil Samples for Presence of ail-bearing *Yersinia enterocolitica* by using a 5' Nuclease Fluorogenic PCR (TaqMan) Assay

Laura Byl

United States Department of Agriculture

Irene V. Wesley

United States Department of Agriculture

Follow this and additional works at: http://lib.dr.iastate.edu/swinereports_2001



Part of the [Agriculture Commons](#), and the [Animal Sciences Commons](#)

Extension Number: ASL R1789

Recommended Citation

Byl, Laura and Wesley, Irene V., "Analysis of National Animal Health Monitoring System Tonsil Samples for Presence of ail-bearing *Yersinia enterocolitica* by using a 5' Nuclease Fluorogenic PCR (TaqMan) Assay" (2002). *Swine Research Report, 2001*. 25.

http://lib.dr.iastate.edu/swinereports_2001/25

This report is brought to you for free and open access by the Animal Science Research Reports at Iowa State University Digital Repository. It has been accepted for inclusion in Swine Research Report, 2001 by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Analysis of National Animal Health Monitoring System Tonsil Samples for Presence of ail-bearing *Yersinia enterocolitica* by using a 5' Nuclease Fluorogenic PCR (TaqMan) Assay

Abstract

The fluorogenic 5' nuclease polymerase chain reaction (TaqMan) assay, which was developed in this laboratory, was further optimized and used to screen hogs for *Yersinia enterocolitica* (4). Tonsil samples (n = 1,647) collected during the National Animal Health Monitoring System (NAHMS) Swine 2001 program were tested for the presence of the ail gene, which is present in virulent *Y. enterocolitica*. Tonsil swabs were collected on farms (n = 101) located in Arkansas, Colorado, Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Carolina, Ohio, Oklahoma, Pennsylvania, South Dakota, Texas and Wisconsin. Of the 1,647 tonsil samples, 11.4% (n = 188) were positive.

Keywords

ASL R1789

Disciplines

Agriculture | Animal Sciences

Analysis of National Animal Health Monitoring System Tonsil Samples for Presence of *ail*-bearing *Yersinia enterocolitica* by using a 5' Nuclease Fluorogenic PCR (TaqMan) Assay

Laura Byl and Irene V. Wesley
National Animal Disease Center, Ames, IA

ASL-R1789

Summary and Implications

The fluorogenic 5' nuclease polymerase chain reaction (TaqMan) assay, which was developed in this laboratory, was further optimized and used to screen hogs for *Yersinia enterocolitica* (4). Tonsil samples (n = 1,647) collected during the National Animal Health Monitoring System (NAHMS) Swine 2001 program were tested for the presence of the *ail* gene, which is present in virulent *Y. enterocolitica*. Tonsil swabs were collected on farms (n = 101) located in Arkansas, Colorado, Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Carolina, Ohio, Oklahoma, Pennsylvania, South Dakota, Texas and Wisconsin. Of the 1,647 tonsil samples, 11.4% (n = 188) were positive.

Introduction

Y. enterocolitica is a major human foodborne pathogen which accounts for ~87,000 cases each year in the United States. Yersiniosis is reported frequently in cooler climates (1,5). The most prevalent clinical signs, especially in children, are abdominal pain, fever, diarrhea, nausea, and vomiting (3). Post-infection symptoms include erythema nodosum and reactive arthritis. Swine are an important animal reservoir of *Y. enterocolitica* strains, including serotypes 0:3 and 0:9, which are pathogenic in humans (1,2,10,11). In one study, virulent strains of *Y. enterocolitica* 0:3 were found in the oral cavity and the gastrointestinal tract of pigs (10). In the United States, serotypes 0:8 and 0:5,27 predominate (1,2,3,5,10,11).

Two *Y. enterocolitica* genes, *inv* and *ail*, play a major role in the ability of this microorganism to enter mammalian cells (9,11).

The fluorogenic polymerase chain reaction (TaqMan) assay is a PCR-based protocol that bypasses post-PCR processing for the detection of the PCR product. The 5' PCR assay uses the 5' → 3' activity of *Thermus aquaticus* DNA polymerase (11) and requires a fluorogenic probe. TaqMan fluorogenic probes consist of an oligonucleotide with reporter dye at the 5' end and a quencher dye at the 3' end. When the probe is intact, the quencher dye suppresses the reporter (6). If the target DNA sequence is present, the probe anneals downstream from one of the primer sites and it is cleaved by the DNA polymerase as the primer is extended. This cleavage

liberates the reporter dye from the quencher dye. The liberated reporter dye increases fluorescence (6,7,11). The increase in fluorogenic signal is a direct consequence of the successful PCR amplification of specific DNA targets (11).

Materials and Methods

Sample Enrichment Procedure. Copan tonsil swabs were collected from hogs (n = 1,647). Swabs were enriched in 10 ml ITC media (2 days at room temperature) after which 100 µl of ITC was streaked to CIN (2 days at room temperature).

DNA Extraction Procedure. DNA extractions were prepared from all the ITC samples as follows. A 1 ml aliquot from the ITC media was centrifuged (2 minutes, 10,000 x g). The supernatant was carefully removed and the remaining pellet was resuspended in 200 µl of PrepMan sample preparation reagent (PE Applied Biosystems, Foster City, CA). The tubes were vortexed to resuspend the pellet and then heated (102 °C for 10 minutes). The tubes were centrifuged (2 minutes at 10,000 x g) to pellet cell debris. The supernatant was transferred into a new microcentrifuge tube and was labeled with the kit and sample number. DNA was also extracted from the presumptive *Yersinia* colonies on the CIN plates. DNA extractions were stored at -20 °C until PCR analysis.

Primer and Probes. The probe and primer sequences target the *Y. enterocolitica* *ail* gene (4; Table 1). Oligonucleotide primers and the fluorescently labeled probe were synthesized by Integrated DNA Technologies (Coralville, IA). The probe was labeled at the 5' end with fluorescent reporter dye FAM-6-carboxy-fluorescein and the 3' end with the quencher dye TAMRA-6-carboxy-tetramethyl-rhodamine.

Fluorogenic 5' Nuclease PCR Conditions. The PCR conditions were described previously (4). Each PCR mixture contained 3.5 mM MgCl₂, 0.2 mM of each deoxynucleoside triphosphate, 1.25 U of AmpliTaq Gold DNA polymerase (PE Applied Biosystems), and 1 X GeneAmp PCR Gold Buffer (150mM Tris-HCl, 500 mM KCl) (PE Applied Biosystems). 5 microliters of DNA template was added to yield a total volume of 50 µl. Concentrations of primers (200 nM) and the probe (25 nM) as originally described (4) yielded inconsistent results. Therefore, primers (50 to 900 nM) were tested in

a matrix format with a known positive DNA template. Optimal concentrations of 300 nM for both the forward and reverse primer were then used to screen the tonsil samples. Probe concentration was also increased from 25 to 200 nM. The thermal cycling conditions were as follows: 95° C for 10 minutes and 35 cycles of 95° C for 15 seconds (denaturation) and 58° C for 1 minute (primer annealing and extension) followed by an indefinite hold at 25° C (4).

Analysis was completed on the ABI 7700 Sequence Detection System. Cycle 33 was set as an arbitrary cut off point. Any sample that crossed the threshold prior to cycle 33 and showed logarithmic amplification was scored as positive. Any sample that crossed the threshold after or on cycle 33 or did not show sufficient amplification was scored as negative (Fig 1).

Results and Discussion

The *ail* gene of pathogenic *Y. enterocolitica* was detected in 11.1% of ITC samples (124 of 1,120) and in 12.1% (64 of 527) of samples plated to CIN (Fig 1).

The fluorogenic PCR assay detected the *ail*-bearing strain of *Y. enterocolitica* in tonsil swabs of normal market-weight hogs. This is evidenced by the 11.1% positive ITC samples and 12.1% positive CIN samples. Swine are known carriers of *Y. enterocolitica*. This study confirmed that virulent species of *Y. enterocolitica* harboring the *ail* gene are present in tonsil swabs of United States swine.

References

1. Cover, T.L. and R.C. Aber. 1989. *Yersinia enterocolitica*. N. Engl. J. Med. 321: 16 - 24.
2. Doyle, M.P., Hugdahl M.B. and Taylor S.L. 1981. Isolation of virulent *Yersinia enterocolitica* from porcine tongues. Appl. Environ. Microbiol. 42: 661 - 666.
3. Funk, J.A., Troutt H.F., Isaacson R.E. and Fossler C.P. 1998. Prevalence of pathogenic *Yersinia enterocolitica* in groups of swine at slaughter. J. Food Prot. 61: 677 - 682.
4. Jourdan, A.D., Johnson S.C. and Wesley I.V. 2000. Development of a fluorogenic 5' nuclease PCR assay for detection of the *ail* gene of pathogenic *Yersinia enterocolitica*. Appl. Environ. Microbiol. 66: 3750 - 3755.
5. Kapperud, G. "Yersinia Enterocolitica Infection." Handbook of Zoonoses, 2nd Ed. Boca Raton, FL: CRC Press, 1994. 343 - 353.
6. Livak, K., Marmaro J., and Flood S. 1995. Guidelines for designing TaqMan fluorogenic probes for 5' nuclease assays. ABI PRISM Sequence Detection Systems Users Bulletin.
7. Livak, K.J. Quantitation of DNA/RNA Using Real-time PCR Detection. Applied Biosystems. May 8, 2001
<http://www.appliedbiosystems.com/molecularbiology/about/pcr/sds/white.html#fn1>
8. Miller, V.L., Farmer III J.J., Hill W.E. and Falkow S. 1989. The *ail* Locus is found uniquely in *Yersinia enterocolitica* serotypes commonly associated with disease. Infect. Immun. 57: 121 - 131.
9. Pierson, D.E. and Falkow S. 1993. The *ail* gene of *Yersinia enterocolitica* has a role in the ability of the organism to survive serum killing. Infect. and Immun. 61: 1846 - 1852.
10. Thibodeau, V., Frost E.H., Chenier S., and Quessy S. 1999. Presence of *Yersinia enterocolitica* in tissues of orally-inoculated pigs and the tonsils and feces of pigs at slaughter. Can. J. Vet. Res. 63: 96 - 100.
11. Vishnubhatla, A., Fung D.Y.C., Oberst R.D., Hays M.P., Nagaraja T.G. and Flood S.J.A. 2000. Rapid 5' nuclease (TaqMan) assay for detection of virulent strains of *Yersinia enterocolitica*. Appl. Environ. Microbiol. 66: 4131 - 4135.

Table 1. List of primers and probe used in the assay.

Primer/probe	Annealing site	Sequence (5' → 3')
TM1-F	746 to 772	GGT CAT GGT GAT GTT GAT TAC TAT TAC
TM1-R	818 to 836	CGG CCC CCA GTA ATC CAT A
TM1-P	785 to 814	CCA TCT TTC CGC ATC AAC GAA TAT CTT AGC